

Derivation of the universal decay cascade distribution

Vyacheslavs Kashcheyevs

Faculty of Physics and Mathematics, University of Latvia, Riga LV-1002, Latvia

A detailed derivation of the decay cascade probability distribution stated in Eqs. (4)–(6) and (11) of Phys. Rev. Lett. **104**, 186805 (2010) [arXiv:0901.4102] by Kashcheyevs and Kaestner is provided. Recurrence relations are solved explicitly and connections between solutions in different limits are demonstrated.

I. MATHEMATICAL DEFINITION OF THE DECAY CASCADE MODEL [2]

The probability distribution $P_n(t)$ is governed by the equation

$$\frac{dP_n(t)}{dt} = -\Gamma_n(t)P_n(t) + \Gamma_{n+1}(t)P_{n+1}(t) \quad (\text{KK-1})$$

for $n \geq 0$ and with $\Gamma_0 \equiv 0$. Normalization and initial conditions are

$$\sum_{n=0}^N P_n(t) = 1, \quad (1)$$

$$P_n(t_0) = \begin{cases} 1, & n = N \\ 0, & n \neq N \end{cases}. \quad (\text{KK-2})$$

Equation (KK-1) is a general kinetic equation for a birth-death Markov process [1] for time- and population-size-dependent rates. Here we will be interested in the asymptotic values of P_n as the transition rates Γ_n gradually decrease to zero as function of time.

The original publication [2] discussing solutions to the above model in the context of dynamic quantum dot initialization is denoted KK; equations marked here as KK-1, KK-2 etc. match the equations in KK with the corresponding numbers.

II. IMPLICIT EXACT SOLUTION

In KK, the following exact iterative solution, valid for $t > t_0$, is presented

$$P_n(t) = \int_{t_0}^t e^{-\int_{t'}^t \Gamma_n(\tau) d\tau} \Gamma_{n+1}(t') P_{n+1}(t') dt', \quad (\text{KK-3})$$

$$P_{N+1}(t) = \delta(t - t_0) / \Gamma_{N+1}(t_0). \quad (2)$$

Condition (2) is introduced formally, in order for the general formula (KK-3) to accommodate the initial condition (KK-2); the delta functions are regularised as $\int_{t_0}^t \delta(t - t_0) = 1$ for $t > t_0$.

Equation (KK-3) is just the standard solution of a single linear first order differential equation (KK-1) for an unknown function $P_n(t)$ with $P_{n+1}(t)$ treated as known. One can verify that (KK-3) solves (KK-1) by direct substitution:

$$\begin{aligned} \frac{dP_n(t)}{dt} &= \underbrace{e^{-\int_t^t \Gamma_n(\tau) d\tau}}_{=1} \Gamma_{n+1}(t) P_{n+1}(t) + \int_{t_0}^t \frac{d}{dt} \left[e^{-\int_{t'}^t \Gamma_n(\tau) d\tau} \right] \Gamma_{n+1}(t') P_{n+1}(t') dt' \\ &= \Gamma_{n+1}(t) P_{n+1}(t) + \int_{t_0}^t e^{-\int_{t'}^t \Gamma_n(\tau) d\tau} \underbrace{\frac{d}{dt} \left[-\int_{t'}^t \Gamma_n(\tau) d\tau \right]}_{=-\Gamma_n(t)} \Gamma_{n+1}(t') P_{n+1}(t') dt' \\ &= \Gamma_{n+1}(t) P_{n+1}(t) - \Gamma_n(t) P_n(t). \end{aligned}$$

III. EXPLICIT SOLUTION FOR TIME-INDEPENDENT RATE RATIO $\Gamma_n(t)/\Gamma_{n-1}(t) = \text{const}$

The first solution described in KK corresponds to the case of

$$\Gamma_n(t) = \frac{X_n}{X_1} \Gamma_1(t), \quad (3)$$

with $X_n \equiv \exp \sum_{k=1}^n \delta_k$ being time-independent constants.

For the rates obeying the condition (3), the general solution can be constructed in the following form ¹

$$P_n(t) = \sum_{k=n}^N R_{nk} e^{-\int_{t_0}^t \Gamma_k(t') dt'}, \quad (4)$$

with constant coefficients R_{nk} that need to be determined.

A. Derivation of R_{nk}

For $n = N$, the initial conditions $P_N(t_0) = 1$, $P_{N+1}(t_0) = 0$ and equation (4) for $P_N(t)$ give

$$P_N(t) = e^{-\int_{t_0}^t \Gamma_N(t') dt'} \implies R_{NN} = 1. \quad (5)$$

The initial condition $P_n(t_0) = 0$ for $n < N$, applied to (4), implies:

$$\sum_{k=n}^N R_{nk} = 0, \quad n < N. \quad (6)$$

For $n < N$ the substituting (4) into the differential equation (4) gives

$$\frac{dP_n(t)}{dt} = - \sum_{k=n}^N R_{nk} \Gamma_k(t) e^{-\int_{t_0}^t \Gamma_k(t') dt'} \quad (7)$$

$$-\Gamma_n(t)P_n(t) + \Gamma_{n+1}(t)P_{n+1}(t) = -\Gamma_n(t) \sum_{k=n}^N R_{nk} e^{-\int_{t_0}^t \Gamma_k(t') dt'} + \Gamma_{n+1}(t) \sum_{k=n+1}^N R_{n+1,k} e^{-\int_{t_0}^t \Gamma_k(t') dt'} \quad (8)$$

Now we invoke the condition (3) which allows us to equate the coefficients of $e^{-\int_{t_0}^t \Gamma_k(t') dt'}$ between (7) and (8):

$$X_k R_{nk} = X_n R_{nk} + X_{n+1} R_{n+1,k}, \quad n < k \leq N \quad (9)$$

$$R_{n-1,k} = \frac{X_n}{X_k - X_{n-1}} R_{n,k}, \quad n \leq k \leq N \quad (9')$$

Equations (6) and (9) give the sought-after recurrence relations:

$$R_{nk} = R_{kk} \prod_{m=n+1}^k \frac{X_m}{X_k - X_{m-1}}, \quad n < k \leq N \quad (\text{KK-5}')$$

$$R_{kk} = - \sum_{m=k+1}^N R_{km}. \quad k < N \quad (\text{KK-6}')$$

Equations (KK-5') and (KK-6') together with (5) are equivalent to Eqs. (5) and (6) of KK with $C_k \equiv R_{kk}$ and $Q_{kn} \equiv R_{kn}/R_{kk}$.

¹ This form was inspired by studying explicit solutions for $\Gamma_n(t) \sim e^{-t}$ and $N = 1, 2, 3$ with the means of a computer algebra system **Mathematica**.

B. Explicit solution of recurrence relations

The recurrence relations (KK-5') and (KK-6') admit the following explicit solution:

$$R_{nk} = \prod_{m=n+1}^N X_m \prod_{\substack{m=n \\ m \neq k}}^N \frac{1}{X_k - X_m}. \quad (10)$$

The solution (10) can be obtained for finite n and k by means of computer algebra, and proven in general form by induction.

Since integration over time preserves the condition (3), we can choose

$$X_n = \int_{t_0}^t \Gamma_n(t) dt \quad (11)$$

and write down the solution explicitly:

$$P_n(t) = \sum_{k=n}^N e^{-X_k} \prod_{m=n+1}^N X_m \prod_{\substack{m=n \\ m \neq k}}^N \frac{1}{X_k - X_m}. \quad (12)$$

The solution (12) agrees precisely with the solution for $N = 3$ and $\Gamma_n = \text{const}$ obtained by Miyamoto *et al.* [3].

IV. SOLUTION IN THE LIMIT OF TIME-SCALE SEPARATION BETWEEN CASCADE STEPS

A more general solution that does not rely on condition (3) is derived in KK in the limit of decay time-scale separation between consecutive steps of the cascade. This is motivated as follows: $P_{n+1}(t)$ stops changing appreciably over a timescale on the order of $\Gamma_{n+1}^{-1}(t)$ during which $P_n(t)$ changes only due to probability flux from state $(n+1)$, with negligible decay down the cascade to state $(n-1)$. This condition corresponds to $\Gamma_{n+1}(t) \gg \Gamma_n(t)$ during the relevant time interval.

The mathematical part of the derivation proceeds as follows:

1. Summing equations (KK-3) for all dP_m/dt with $N \geq m > n$ gives

$$\Gamma_{n+1}(t)P_{n+1}(t) = -\frac{d}{dt} \sum_{m>n} P_m(t). \quad (13)$$

2. On the “slow” time-scale controlled by $\Gamma_n(t)$, the function $\sum_{m>n} P_m(t)$ is changing rapidly from 1 to its asymptotic value $\sum_{m>n} P_m(t \rightarrow \infty)$ and hence can be approximated by a step function in time. ($t \rightarrow \infty$ corresponds to the time when the decay transitions no longer take place).
3. The derivative of a step function is proportional to a delta function and thus the exact integral (KK-3) can be approximated as follows:

$$P_n(t) = \int_{t_0}^t e^{-\int_{t'}^t \Gamma_n(\tau) d\tau} \underbrace{\Gamma_{n+1}(t') P_{n+1}(t')}_{\approx \delta(t_0 - t') [1 - \sum_{m>n} P_m(t \rightarrow \infty)]} dt' \approx \underbrace{e^{-\int_{t_0}^t \Gamma_n(\tau) d\tau}}_{e^{-X_n}} [1 - \sum_{m>n} P_m(t \rightarrow \infty)]. \quad (14)$$

Equation (14) essentially states that the decay of all previous states (higher than n) provides an initial condition for the decay of the n -th state. Based on (14), the condition on the final probabilities $P_n(t \rightarrow \infty) \equiv P_n$ stated in KK is formulated:

$$P_n = e^{-X_n} \left(1 - \sum_{m=n+1}^N P_m \right). \quad (15)$$

4. Equation (15) is solved by expressing P_{n-1} in terms of P_n . Using (15) for P_n we can express $\sum_{m>n} P_m = 1 - e^{X_n} P_n$ and

$$\sum_{m>n-1} P_m = 1 + (1 - e^{X_n}) P_n. \quad (16)$$

Substituting the sum (16) into (15) for P_{n-1} , we get the derised recurrence relation

$$P_{n-1} = e^{-X_{n-1}}(e^{X_n} - 1)P_n. \quad (17)$$

Starting from $P_N = e^{-X_N}$ and iterating (17) gives

$$\underbrace{\underbrace{\underbrace{e^{-X_N} \times (e^{X_N} - 1)}_{P_N} e^{-X_{N-1}} \times (e^{X_{N-1}} - 1)}_{P_{N-1}} e^{-X_{N-2}} \times \dots}_{P_{N-2}}, \quad (18)$$

wherefrom the general form is easy to infer

$$P_n = e^{-X_n} \prod_{m=n+1}^N (1 - e^{-X_m}). \quad (\text{KK-11})$$

Equation (KK-11) has also been applied to a generalised decay cascade scenario by Fricke *et al.* [4]. They consider the onset of decay steps at different times, $t_0 < \dots < t_{n+1}^b < t_n^b < \dots$, which in the notation of KK corresponds to

$$\Gamma_n(t) = \tilde{\Gamma}_n(t) \Theta(t - t_n^b), \quad (19)$$

where $\Theta(x)$ is a unit step function and $\tilde{\Gamma}_n(t)$ are smooth functions that decay to zero as for $t \rightarrow \infty$. The condition $X_n = \int_{t_0}^{\infty} \Gamma_n(t) dt = \int_{t_n^b}^{\infty} \tilde{\Gamma}_n(t) dt \gg X_{n-1}$ justifies the sequential cascade approximation (14) and hence the probability distribution (KK-11).

-
- [1] C. Gardiner, *Stochastic Methods: A Handbook for the Natural and Social Sciences* (Springer Verlag, 2009), 4th ed.
 - [2] V. Kashcheyevs and B. Kaestner, Phys. Rev. Lett. **104**, 186805 (2010).
 - [3] S. Miyamoto, K. Nishiguchi, Y. Ono, K. M. Itoh, and A. Fujiwara, Appl. Phys. Lett. **93**, 222103 (2008).
 - [4] L. Fricke, M. Wulf, B. Kaestner, V. Kashcheyevs, J. Timoshenko, P. Nazarov, F. Hohls, P. Mirovsky, B. Mackrodt, R. Dolata, et al., Phys. Rev. Lett. **110**, 126803 (2013).